arXiv:1211.4959v1 [math.AP] 21 Nov 2012

APPROXIMATION AND EQUIDISTRIBUTION OF PHASE SHIFTS: SPHERICAL SYMMETRY

KIRIL DATCHEV, JESSE GELL-REDMAN, ANDREW HASSELL, AND PETER HUMPHRIES

Abstract. Consider a semiclassical Hamiltonian

$$H_{V,h} := h^2\Delta + V - E$$

where $h > 0$ is a semiclassical parameter, Δ is the positive Laplacian on $\mathbb{R}^d$, $V \in C_c^\infty(\mathbb{R}^d)$, i.e. V is a smooth, compactly supported potential function which is central, that is, depends only on $|x|$ and $E > 0$ is an energy level. In this setting the scattering matrix $S_h(E)$ can be defined at any positive energy E; it is a unitary operator on $L^2(\mathbb{S}^{d-1})$, hence with spectrum lying on the unit circle; moreover, the spectrum is discrete except at 1.

We show under certain additional assumptions on the potential that the eigenvalues of $S_h(E)$ can be divided into two classes: a finite number $\sim c_d(R\sqrt{E}/h)^{d-1}$, as $h \to 0$, where $B(0,R)$ is the convex hull of the support of the potential, that equidistribute around the unit circle, and the remainder that are all very close to 1. Semiclassically, these are related to the rays that meet the support of, and hence are scattered by, the potential, and those that do not meet the support of the potential, respectively.

A similar property is shown for the obstacle problem in the case that the obstacle is the ball of radius R.

1. Introduction

In this paper we consider the scattering matrix for a semiclassical potential scattering problem with spherical symmetry on $\mathbb{R}^d$, $d \geq 2$. Let V be a smooth, compactly supported potential function which is central, i.e. $V(x)$ depends only on $|x|$. We consider the Hamiltonian

$$H_{V,h} := h^2\Delta + V - E \tag{1.1}$$

where $\Delta = -\sum_{i=1}^d \partial_i^2$ is the positive Laplacian on $\mathbb{R}^d$, $E > 0$ is a positive constant (energy) and $h > 0$ is a semiclassical parameter.

The scattering matrix $S_h(E)$ for this Hamiltonian is defined in terms of the asymptotics of generalized eigenfunctions of $H_{V,h}$. For each function $q_{in} \in C^\infty(\mathbb{S}^{d-1})$, there is a unique solution to $H_{V,h}u = 0$ of the form

$$u = r^{-(d-1)/2}\left(e^{-i\sqrt{E}r/h}q_{\text{in}}(\omega) + e^{+i\sqrt{E}r/h}q_{\text{out}}(-\omega)\right) + O(r^{-(d+1)/2}), \tag{1.2}$$

as $r \to \infty$, see e.g. [15]. Here $q_{\text{out}} \in C^\infty(\mathbb{S}^{d-1})$. The map $q_{\text{in}} \mapsto e^{i\pi(d-1)/2}q_{\text{out}}$ is by definition the scattering matrix $S_h(E)$. The factor $e^{i\pi(d-1)/2}$ is chosen so that this 'stationary' definition agrees with time-dependent definitions (see e.g. [17] or [18]), and is such that the

---

The first author was partially supported by an NSF postdoctoral fellowship. The authors would like to thank Hamid Hezari, Volker Schlue and Steve Zelditch for very helpful conversations. The third author acknowledges the support of the Australian Research Council through a Future Fellowship FT0990895 and Discovery Grant DP1095448.

scattering matrix for the potential $V \equiv 0$ is the identity. It is standard that the scattering matrix $S_h(E)$ is, for every $h > 0$, a unitary operator on $L^2(\mathbb{S}^{d-1})$, and that $S_h(E) - \mathrm{Id}$ is compact. It follows that the spectrum lies on the unit circle, consists only of eigenvalues, and is discrete except at 1. It is therefore possible to count the number of eigenvalues of $S_h(E)$ in any closed interval of the unit circle not containing 1. In fact, semiclassically (i.e. as $h \to 0$) we are able to separate the spectrum of $S_h(E)$ into two parts. One is associated to the rays that meet the support of the potential; to leading order in h there are $c_d(R\sqrt{E}/h)^{d-1}$ of these, $c_d = 2/(d-1)!$, and the other part is associated to the rays that do not meet the support of the potential. Those eigenvalues corresponding to rays that do not meet the support are close to 1, as one should expect, since the eigenvalues of the zero potential are all 1 — see Proposition 1.3. The other eigenvalues are affected by the potential, and we can ask whether these 'nontrivial' eigenvalues are asymptotically equidistributed on the unit circle. Indeed Steve Zelditch posed this question to one of the authors several years ago.

Before stating the main result, we discuss further the scattering matrix in the case of central potentials. In this case the eigenfunctions of the scattering matrix are spherical harmonics and the Helmholtz equation reduces to an ODE in the radial variable. The generalized eigenfunctions then take the form $u = r^{-(d-2)/2} f(r) Y_l^m$, where

$$\left(-\partial_r^2 - \frac{1}{r}\partial_r - \frac{(l + (d-2)/2)^2}{r^2} + \frac{V(r) - E}{h^2} \right) f = 0 \tag{1.3}$$

and $f(r) = H^{(1)}_{l+(d-2)/2}(r\sqrt{E}/h) + c(l) H^{(2)}_{l+(d-2)/2}(r\sqrt{E}/h)$ for $r > R$, where the $H^{(i)}_\nu$ are the standard Hankel functions, [1]. With our normalization, $S_h Y_l^m = c(l) Y_l^m$. In particular, *the eigenvalue of* Y_l^m *is independent of* m. We write the eigenvalue corresponding to Y_l^m in the form $e^{i\beta_{l,h}}$. The quantities $\beta_{l,h}/2$ are frequently referred to as 'phase shifts.' See e.g. [17] for a review of these facts.

We now discuss the conditions we on the potentials in the main theorem. Both conditions are dynamical conditions, i.e. conditions on the Hamiltonian dynamical system determined by the symbol of $H_{V,h}$. As usual in microlocal analysis we refer to the classical trajectories of this system as bicharacteristics. We first define the interaction region

$$\mathcal{R} := \{x : V(|y|) < E \text{ for all } |y| > |x|\}\,. \tag{1.4}$$

This is the region of x-space accessible by bicharacteristics coming from infinity. Notice that for central potentials this region takes the form

$$\mathcal{R} = \{|x| \geq r_0\} \text{ where } r_0 = \inf_{r \geq 0}\{s > r \Rightarrow V(s) < E\}. \tag{1.5}$$

The first condition is

$$V \text{ is nontrapping at energy } E \text{ in the interaction region.} \tag{1.6}$$

That is, x tends to infinity along every bicharacteristic in $\mathcal{R}$ both forwards and backwards in time.

The second concerns the *scattering angle.* Let R be such that $B(0, R)$ is the smallest ball containing the support of V, i.e.

$$B(0, R) = \mathrm{chsupp} V, \text{ the convex hull of the support of V.} \tag{1.7}$$

We recall (see Section 2 for further details) that for a central potential, the scattering angle $\Sigma(\alpha)$ is a function only of the angular momentum α and measures the difference between the incident and final directions of the trajectory (which are well-defined, since the motion is free for $|x| > R$ where $B(0, R)$ is the support of the potential) — see [17]. The scattering angle is zero for all trajectories with $\alpha > R$. Our second condition is that

$$\text{The zeroes of } \Sigma'(\alpha) = \frac{d}{d\alpha}\Sigma(\alpha) \text{ are finite in } [0, R). \tag{1.8}$$

Then our main result is

Theorem 1.1. *Let R be as in* (1.7)*, and assume that $V \in C_c^\infty(\mathbb{R}^d)$ be central and satisfies conditions* (1.6) *and* (1.8)*. Then as $h \downarrow 0$, we consider the eigenvalues $e^{i\beta_{l,h}}$ for which $l \leq R/h$, counted with multiplicity $p_d(l) = \dim\ker(\Delta_{\mathbb{S}^{d-1}} - l(l+d-2))$. There are $2(R\sqrt{E}/h)^{d-1}/(d-1)! + O(h^{-(d-2)})$ of these, and they equidistribute around the unit circle, meaning that*

$$\sup_{0\leq\phi_0<\phi_1\leq 2\pi}\left|\frac{N(\phi_0,\phi_1)}{2(R\sqrt{E}/h)^{d-1}/(d-1)!} - \frac{\phi_1-\phi_0}{2\pi}\right| \to 0 \text{ as } h \downarrow 0, \tag{1.9}$$

where $N(\phi_0, \phi_1)$ is the number of $\beta_{l,h}$ with $l \leq R/h$ and $\phi_0 \leq \beta_{l,h} \leq \phi_1$ (mod 2π), counted with multiplicity.

Remark 1.2. Many potentials satisfy conditions (1.6) and (1.8). For example, take any smooth compactly supported function $W(r)$ such that $W(r) = 0$ for $r \geq R$, $W(r) > 0$ for $0 \leq r < R$ and $W''(r)$ is positive and monotone decreasing in some nonempty interval $[R - \epsilon, R)$. Then $cW(r)$ satisfies the conditions for sufficiently large c. An explicit example is

$$W(r) = \begin{cases} e^{1/(r^2-R^2)}, & r < R \\ 0, & r \geq R. \end{cases}$$

More generally, in Section 5 we show that a potential satisfying $V' \leq 0$ and $V' + rV'' > 0$ on the interaction region $\mathcal{R}$ also satisfies conditions (1.6) and (1.8). This condition is actually much stronger than (1.8): it implies that $\Sigma'(\alpha)$ has no zeroes at all on $[0, R)$.

We also show

Proposition 1.3. *Let $\kappa \in (0, 1)$. The eigenvalues $e^{i\beta_{l,h}}$ for $l \geq (R + h^\kappa)/h$ satisfy*

$$\left|e^{i\beta_{l,h}} - 1\right| = O(h^\infty), \quad h \to 0.$$

Here and below, $O(h^\infty)$ denotes a quantity that is bounded by $C_N h^N$ for all N and some $C_N > 0$.

Note that the number of eigenvalues not covered by the Theorem and Proposition above is $o(h^{1-d})$, and hence cannot affect the equidistribution properties. Hence we get

Corollary 1.4. *Suppose that V satisfies conditions* (1.6) *and* (1.8)*. Then for each $\epsilon > 0$, we have*

$$\sup_{\epsilon\leq\phi_0<\phi_1\leq 2\pi-\epsilon}\left|\frac{\tilde{N}(\phi_0,\phi_1)}{2(R\sqrt{E}/h)^{d-1}/(d-1)!} - \frac{\phi_1-\phi_0}{2\pi}\right| \to 0 \text{ as } h \downarrow 0, \tag{1.10}$$

where $\tilde{N}(\phi_0, \phi_1)$ is the number of $\beta_{l,h}$ (with no condition on l), counted with multiplicity, satisfying $\phi_0 < \beta_{l,h} < \phi_1$ (mod 2π).

Results directly analogous to those for semiclassical potentials are also true in the case of scattering by a disk of radius R centered at the origin. The scattering matrix in this case can be defined similarly; given any function $q_{\text{in}} \in C^\infty(\mathbb{S}^{d-1})$, there is a unique solution u to the equation $\left(\Delta - k^2\right) u = 0$ such that[1]

$$\begin{aligned} u &= r^{-(d-1)/2}\left(e^{-ikr} q_{\text{in}} + e^{+ikr} q_{\text{out}}\right) + O(r)^{-(d+1)/2}, \quad r \to \infty \\ u|_{|x|=R} &\equiv 0. \end{aligned} \tag{1.11}$$

The scattering matrix S_k is again defined $q_{\text{in}} \mapsto e^{i\pi(d-1)/2} q_{\text{out}}$, and the standard facts about the operator S_h also hold for S_k. As above, the spherical harmonics diagonalize the scattering matrix. We write $S_k Y_l^m = e^{ix_{l,k}} Y_l^m$. We will prove

Theorem 1.5. *As $k \to \infty$, the eigenvalues $e^{ix_{l,k}}$ for which $l \leq Rk$, counted with multiplicity $p_d(l) = \dim\ker\left(\Delta_{\mathbb{S}^{d-1}} - l(l+d-2)\right)$, equidistribute around $\mathbb{S}^1$ in the sense of the Main Theorem. In fact, we have the stronger statement*

$$\sup_{0 \leq \phi_0 < \phi_1 \leq 2\pi} \left| \frac{N(\phi_0, \phi_1)}{2(Rk)^{d-1}/(d-1)!} - \frac{\phi_1 - \phi_0}{2\pi} \right| = O(k^{-1/3}) \text{ as } k \to \infty. \tag{1.12}$$

As far as we are aware, the present paper is the first in the mathematical literature to deal with the question of the equidistribution of phase shifts over the unit circle. However, there are a number of previous studies of high-energy or semiclassical asymptotics of eigenvalues of the scattering matrix. The relation between the sojourn time and high-frequency asymptotics of the scattering matrix was observed in classical papers by Guillemin and Majda. Melrose and Zworski [16] showed that for fixed $h > 0$ the absolute scattering matrix for a Schrödinger operator on a scattering, or asymptotically conic, manifold is an FIO associated to the geodesic flow on the manifold at infinity for time π. Alexandrova [2] studied the scattering matrix for a nontrapping semiclassical Schrödinger operator, and showed that localized to finite frequency, it is a semiclassical FIO associated to the limiting Hamilton flow relation at infinity, which includes the behavior of the Hamilton flow in compact sets. A more global description was given in Hassell-Wunsch [9] where the semiclassical asymptotics of the scattering matrix were unified with the singularities of the scattering matrix at fixed frequency (i.e. the Melrose-Zworski result [16]). These results are explained in Sections 2 and 3 below.

Asymptotics of phase shifts, i.e. the eigenvalues of the scattering matrix, were analysed by Birman-Yafaev [3, 4, 5, 6], Yafaev [19] and more recently Bulger-Pushnitski [7]. This was done in the context of the scattering theory for a fixed potential. In this case, the scattering matrix $S(\lambda)$ tends in operator norm to the identity as $\lambda \to \infty$ so the phase shifts tend to zero uniformly. The asymptotics of the phase shifts for a fixed energy, and also the high-energy asymptotics, were analyzed.

In [21], Zelditch and Zworski analyzed the pair correlation function for eigenvalues of the scattering matrix associated to a rotationally invariant surface with a conic singularity and cylindrical ends. They showed that a full measure set of a 2-parameter family of such surfaces obeyed Poisson statistics, as conjectured by Smilansky, motivated by the Berry-Tabor conjecture for completely integrable Hamiltonian systems.

In a slightly different setting Zelditch [20] analyzed quantized contact transformations, which are families of unitary maps on finite dimensional spaces with dimension $N \to \infty$. He proved under the assumption that the set of periodic points of the transformation has measure zero, that the eigenvalues of these unitary operators becomes equidistributed as $N \to \infty$. After reading a draft of the current paper, Zelditch pointed out to the authors that a similar strategy could be used in the context of semiclassical potential scattering to prove equidistribution. In fact, this strategy is likely to be a more direct approach to proving equidistribution than the one we employ here. On the other hand, our approach has several advantages: it also gives approximations to the individual phase shifts, up to an $O(h)$ error (see Proposition 3.1), and in addition appears to be a better method for obtaining a rate of equidistribution, as in Theorem 1.5 above.

---

[1] Here we prefer to use non-semiclassical notation where the energy level is k^2, as is traditional in obstacle scattering literature. The variable k here corresponds to $1/h$ above, when the energy level $E = 1$.

2. Dynamics

We now review some standard material on Hamiltonian dynamics for central potentials. Consider first the case the dimension $d = 2$.

The classical Hamiltonian corresponding to our quantum system is

$$|\xi|^2 + V(r) - E$$

or in polar coordinates, using (r, φ) and dual coordinates (ρ, η),

$$H = \rho^2 + \frac{\eta^2}{r^2} + V(r) - E.$$

The Hamilton equations of motion are

$$\begin{aligned} \dot{r} &= 2\rho \\ \dot{\varphi} &= 2\frac{\eta}{r^2} \\ \dot{\rho} &= -V'(r) + 2\frac{\eta^2}{r^3} \\ \dot{\eta} &= 0. \end{aligned} \tag{2.1}$$

The invariance of the Hamiltonian under rotations is reflected in the conservation of angular momentum $\eta = 2r^2\dot{\varphi}$. For a given bicharacteristic, this is the minimum value of r along the free bicharacteristic that agrees with the given one as $t \to -\infty$ (we could just as well take $t \to +\infty$ since it is a conserved quantity).

Notice that in the case of general dimension d, each bicharacteristic lies entirely in a two-dimensional subspace, so the above discussion in fact includes the general case.

The scattering matrix is related to the asymptotic properties of the bicharacteristic flow. Geometrically this information is contained in a submanifold

$$L \subset \mathrm{T}^* \mathbb{S}^{d-1} \times \mathrm{T}^* \mathbb{S}^{d-1} \times \mathbb{R}$$

that we define now. Returning to the case of general dimension d, we identify $\mathbb{S}^{d-1}$ with the unit sphere in $\mathbb{R}^d$ and identify the cotangent space $\mathrm{T}^*_\omega \mathbb{S}^{d-1}$ with the orthogonal hyperplane $\omega^\perp$ to ω. Given ω and $\eta \in \mathrm{T}^*_\omega \mathbb{S}^{d-1}$, take the unique bicharacteristic ray whose projection $x_{\omega,\eta}(t)$ to $\mathbb{R}^d$ is given by $\eta + t\omega$ for $t << 0$. Define (ω', η') by

$$\begin{aligned} \omega'(\omega, \eta) &= \lim_{t\to\infty} x(t)/\,|x(t)| \\ \eta'(\omega, \eta) &= \lim_{t\to\infty} x(t) - \langle x(t), \omega' \rangle \omega' \end{aligned} \tag{2.2}$$

and $\tau(\omega, \eta)$ to be the **sojourn time** or **time delay** for γ; this is by definition the limit

$$\lim_{a\to\infty, a>>0} t_1(a) - t_2(a) - 2a = \tau(\omega, \eta), \tag{2.3}$$



FIGURE 1. Here $x_{\omega,\eta}$ is the classical trajectory equal to $\eta + t\omega$ for $t << 0$. The scattering angle $\Sigma(\eta)$ is the difference between the outgoing direction ω' and the incoming direction ω. Note that $|\eta| = |\eta'|$ by conservation of angular momentum since the potential V is central, $V = V(r)$.

where $t_1(a)$ is the smallest time, t, for which $r(t) = a$ and $t_2(a)$ is the largest. We then define L to be the submanifold

$$L := \{(\omega, \eta, \omega'(\omega, \eta), -\eta'(\omega, \eta), \tau).\} \tag{2.4}$$

As shown in [9], L is a Legendrian submanifold of $\mathrm{T}^* \mathbb{S}^{d-1} \times \mathrm{T}^* \mathbb{S}^{d-1} \times \mathbb{R}$ with respect to the contact form $\chi + \chi' - d\tau$, where χ is the standard contact form on $\mathrm{T}^* \mathbb{S}^{d-1}$, given in any local coordinates x and dual coordinates ξ by $\xi \cdot dx$. Note that the projection of L to $\mathrm{T}^* \mathbb{S}^{d-1} \times \mathrm{T}^* \mathbb{S}^{d-1}$ is Lagrangian with respect to the standard symplectic form. Indeed it is the graph of a symplectic transformation $(\omega, \eta) \mapsto (\omega', \eta')$, and the scattering matrix is a semiclassical Fourier Integral Operator associated to this symplectic graph [2], [9]. The sojourn time, however, carries extra information and is directly related to high-energy scattering asymptotics as observed in [14], [8], [9].

The previous paragraph applies to any potential, central or not. We return to the case of a central potential, for which, as observed above, the dynamics take place in a two-dimensional subspace, so we can assume $d = 2$ without loss of generality. In that case we use the angular variables φ, φ' in dimension $d = 2$ instead of ω, ω' above. Consider a bicharacteristic with angular momentum $\eta \in \mathbb{R}$ and initial direction $\varphi \in \mathbb{S}^1$. The **scattering angle** $\Sigma(\eta)$ is defined to be the angle between the initial and final directions, normalized so that Σ is continuous and $\Sigma(\eta) = 0$ for $\eta > R$, i.e.

$$\Sigma(\eta) = \varphi'(\varphi, \eta) - \varphi, \quad \Sigma(\eta) = 0 \text{ for } \eta > R, \tag{2.5}$$

(see Figure 1) and the sojourn time is defined by

$$T(\eta) = \tau(\varphi, \eta). \tag{2.6}$$

Notice that both Σ and T depend only on η in the central case. The fact that L in (2.4) is Legendrian then implies the following relation between these functions:

$$d\tau = \eta \cdot (d\varphi - d\varphi') \implies \frac{d}{d\eta}T(\eta) = -\eta \frac{d}{d\eta}\Sigma(\eta). \tag{2.7}$$

Remark 2.1. Notice that the ambiguity of Σ modulo 2π is eliminated by our convention that $\Sigma(\eta) = 0$ for $\eta > R$. We point out that by reflection symmetry, we have $\Sigma(\eta) = -\Sigma(-\eta)$ modulo 2π, but it might not be the case that $\Sigma(\eta) = -\Sigma(-\eta)$ on the nose: this will happen if and only if $\Sigma(0) = 0$, which will be the case if and only if the interaction region is the whole of $\mathbb{R}^d$. However, we always have $\Sigma'(\eta) = \Sigma'(-\eta)$, which shows that $T'(\eta)$ is an odd function, and hence $T(\eta)$ is even in η.

3. An asymptotic formula for the eigenvalues of S_h

In this section we find an asymptotic formula for the eigenvalues $e^{i\beta_{l,h}}$ of S_h in terms of the scattering angle Σ.

Proposition 3.1. *Define the real-valued function* $G(\alpha)$, $\alpha \in \mathbb{R}$, *by*

$$\frac{dG}{d\alpha}(\alpha) = \Sigma(\alpha), \quad G(\alpha) = 0 \text{ for } \alpha \geq R, \tag{3.1}$$

where Σ *is the scattering angle function in* (2.5). *Then we have the following estimate on each eigenvalue* $e^{i\beta_{l,h}}$ *of* S_h*:*

(i) If the dimension d *is even, then for any* $\epsilon > 0$ *there exists* $C = C(\epsilon, d)$ *such that, for all* l *satisfying* $\epsilon \leq lh \leq R$, *we have an estimate*

$$\left| e^{i\beta_{l,h}} - \exp\left\{ \frac{i}{h} \left(G\big((l + \frac{d-2}{2})h\big) \right) \right\} \right| \leq Ch. \tag{3.2}$$

(ii) If the dimension $d > 2$ *is odd, then for any* $\epsilon > 0$ *there exists* $C = C(\epsilon, d)$ *such that* (3.2) *holds whenever* $\alpha = lh \geq \epsilon$ *is distance at least* ϵ *from the set*

$$\left\{ \alpha : \Sigma(\alpha) \in \{\pi k\}_{k \in \mathbb{Z}} \right\}. \tag{3.3}$$

To prove the proposition, we will use the fact, proven in [9], [2], that the integral kernel of S_h is an oscillatory integral associated (in a manner we describe directly) to the Legendre submanifold L in (2.4). To be precise, the Schwartz kernel of S_h can be decomposed following [9, Prop. 15] (with minor changes in notation) as

$$S_h = K_1 + K_2 + K_3,$$

with the K_i as follows.

Fix $R_2 > R_1 > R$. First, K_2 is a pseudodifferential operator of order zero (both in the sense of semiclassical order and differential order), microsupported in $\{|\eta| > R_1\}$, hence taking the form in local coordinates z on $\mathbb{S}^{d-1}$

$$(2\pi h)^{-(d-1)} \int e^{i(z-z')\cdot\zeta/h} b(z,\zeta,h)\, d\zeta$$

for some smooth symbol $b(z,\zeta,h)$ equal to zero for $|\zeta|_{g(z)} < R_1$ where $|\cdot|_{g(z)}$ is the norm on $\mathrm{T}^*_z\,\mathbb{S}^{d-1}$. This reflects the fact that the Legendrian submanifold L in (2.4) is the diagonal relation $\omega = \omega', \eta = -\eta', \tau = 0$ for $|\eta|, |\eta'| > R$, to which pseudodifferential operators are associated. Moreover, K_2 is microlocally equal to the identity for $|\eta| > R_2$, i.e. $b = 1 + O(h^\infty)$ for $|\zeta|_g > R_2$. Indeed, the full symbol (up to $O(h^\infty)$) of the scattering matrix is determined by transport equations along the rays with $|\eta| > R$. Since these transport equations are identical to those for the zero potential, the scattering matrix in this microlocal region is microlocally identical to that for the zero potential, which is the identity operator.

Next, K_1 is a semiclassical Fourier integral operator of semiclassical order 0 with compact microsupport in $\{|\eta| < R_2\}$. That is, K_1 is given by a sum of terms taking the form in local coordinates

$$K_1(\omega,\omega',h) = h^{-(d-1)/2-N/2} \int_{\mathbb{R}^N} e^{i\Phi(\omega,\omega',v)/h} a(\omega,\omega',v,h)dv \tag{3.4}$$

with respect to a suitable phase function Φ and smooth compactly supported function a. Here the phase function parametrizes L locally, meaning

(1) On the set $\operatorname{Crit}\Phi := \{(\omega,\omega',v) : D_v\Phi(\omega,\omega',v) = 0\}$, $D_{\omega,\omega',v}\Phi$ has rank N. This implies that

$$L(\Phi) := (\omega, D_\omega\Phi(\omega,\omega',v), \omega', D_{\omega'}\Phi(\omega,\omega',v), \Phi(\omega,\omega',v)) \tag{3.5}$$

is a smooth submanifold.

(2) $L(\Phi) \cap \operatorname{supp} a = L \cap \operatorname{supp} a$.

By K_1 having compact microsupport in the set $\{|\eta| < R_2\}$, we mean specifically that if $(\omega,\eta,\omega',\eta',\tau) \in L$ has $|\eta| = |\eta|' > R_2$ and $(\omega,\omega',v) \in \operatorname{Crit}(\Phi)$ with $(D_{\omega,\omega'}\Phi(\omega,\omega',v_i),\Phi) = (\eta,\eta',\tau)$, then $a(\omega,\omega',v,h) = O(h^\infty)$ in a neighbourhood of (ω,ω',v).

Finally, K_3 is a kernel in $\dot{C}^\infty(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1} \times [0,h_0))$, i.e. smooth and vanishing to all orders at $h = 0$.

For the proof of Proposition 3.1 we need to know the principal symbol of K_1.

Lemma 3.2. *The Maslov bundle of the canonical relation C of the FIO K_1 is canonically trivial, and with respect to this canonical trivialization, the principal symbol of K_1 is equal to* 1*, as a multiple of the Liouville half-density on C coming from either the left or right projection of C to $\mathrm{T}^*\,\mathbb{S}^{d-1}$. That is to say,*

$$\sigma(K_1)(\omega,\eta,\omega',-\eta') = |d\omega\, d\eta\, d\omega'\, d\eta'|^{1/2}, \tag{3.6}$$

for $(\omega,\eta,\omega',-\eta') \in C$ such that $|\eta| \le R_1$.

Proof. Consider first the Maslov bundle of C. Notice that C is almost the same as L; in fact, it is given by

$$C = \{(\omega,\eta,\omega',-\eta') \mid \exists\,\tau \text{ such that } (\omega,\eta,\omega',\eta',\tau) \in L\}.$$

Since C is a canonical graph (i.e. the graph of a symplectomorphism), associated to the scattering relation as in (2.2), it projects diffeomorphically to $\mathrm{T}^*\,\mathbb{S}^{d-1}$ via both the left and right projections, and the lift of Liouville measure on $\mathrm{T}^*\,\mathbb{S}^{d-1}$ via the left projection agrees with the lift via the right projection (since C is a Lagrangian submanifold of $\mathrm{T}^*\,\mathbb{S}^{d-1}\times \mathrm{T}^*\,\mathbb{S}^{d-1}$ and the Liouville measure can be expressed in terms of the symplectic form $\mathrm{T}^*\,\mathbb{S}^{d-1}$), providing a canonical half-density on C. We also note that the scattering relation is the identity whenever $|\eta| \geq R$ since then the corresponding bicharacteristic is not affected by the potential. Therefore, over this part of C there is a canonical trivialization of the Maslov bundle. Since the Maslov bundle is flat, we can use parallel transport to extend this to a global trivialization: in fact, in the case $d = 2$, the space $\mathrm{T}^*\,\mathbb{S}^1$ retracts to $\mathbb{S}^{d-1}\times\{\eta > R\}$, while for $d \geq 3$, $\mathrm{T}^*\,\mathbb{S}^{d-1}$ is simply connected, hence in either case parallel transport provides an unambiguous trivialization.

We now consider the principal symbol of the scattering matrix. The scattering matrix may be viewed as a 'boundary value' (after removing a vanishing factor and an oscillatory term) of the Poisson operator, as in [9, Section 7.7 and Section 15]. The principal symbol of the scattering matrix is correspondingly derived from the principal symbol of the Poisson operator. The principal symbol of the Poisson operator is real: it solves a real transport equation with initial condition 1. Therefore, the principal symbol of the scattering matrix is real, up to Maslov factors, i.e. it is a real number times an eighth root of unity. On the other hand, unitarity of the scattering matrix shows that the principal symbol lies on the unit circle (as a multiple of the canonical half-density); hence it is an eighth root of unity. Finally, the principal symbol of the scattering matrix is equal to 1 for $|\eta| \geq R$, since here the scattering matrix is microlocally equal to the scattering matrix for the zero potential, which is certainly equal to 1. Since the principal symbol is smooth, is restricted to eighth roots of unity, and is 1 for $|\eta| \geq R$, it follows that the principal symbol is equal to 1 everywhere. ☐

Proof of Proposition 3.1: First we reduce the problem to the cases $d = 2$ and $d = 3$ as follows. Writing $\beta_{l,h,d}$ for the eigenvalue $\beta_{l,h}$ in dimension d, observe that by (1.3),

$$\beta_{l,h,d+2k} = \beta_{l+k,h,d} \text{ for } d \geq 2, k \geq 0. \tag{3.7}$$

It follows that for $d \geq 4$ even, we have $\beta_{l,h,d} = \beta_{l+(d-2)/2,h,2}$ and for $d \geq 5$ odd, we have $\beta_{l,h,d} = \beta_{l+(d-3)/2,h,3}$.

Consider the case dimension $d = 2$. For any smooth function $G : \mathbb{R} \to \mathbb{R}$, the function

$$\Phi(\varphi, \varphi', v) = (\varphi - \varphi')v + G(v), \tag{3.8}$$

locally parametrizes (see (3.5)) the Legendrian

$$L(\Phi) := \big\{(\varphi, \eta, \varphi', \eta', \tau) \mid \eta = v = -\eta', \varphi' - \varphi = \frac{dG}{dv}(v),\ \tau = -v\frac{dG}{dv}(v) + G(v)\big\}.$$

With G as in (3.1), this gives an explicit global parametrization of the Legendrian submanifold L in (2.4) if we take $\varphi \in [0, 2\pi], \varphi' \in \mathbb{R}$. In this case the relation between τ and Σ given by the last equation above is

$$\tau = -\eta\Sigma(\eta) + G(\eta) \implies \frac{d}{d\eta}\tau = -\eta\frac{d}{d\eta}\Sigma(\eta),$$

in agreement with (2.7). Therefore, plugging (3.8) into (3.4), the operator K_1 takes the form

$$K_1(\varphi, \varphi', h) = (2\pi h)^{-1}\int_{\mathbb{R}} e^{i\big((\varphi-\varphi')v+G(v)\big)/h} a(\varphi - \varphi', v, h)\, dv, \quad \varphi \in [0, 2\pi], \varphi' \in \mathbb{R},$$

where a is smooth and supported in $|v| \leq R_2$. Notice that we may assume that a depends only on $(\varphi - \varphi', v, h)$ since the scattering matrix and the phase function both have this property.

Now we obtain an expression for the eigenvalue $e^{i\beta_{l,h}}$ of the scattering matrix S_h on $Y_l = (2\pi)^{-1/2}e^{il\varphi}$ using the pairing

$$e^{i\beta_{l,h}} = \langle S_h Y_l, Y_l\rangle = \langle K_1 Y_l, Y_l\rangle + \langle K_2 Y_l, Y_l\rangle + \langle K_3 Y_l, Y_l\rangle. \tag{3.9}$$

Clearly $\langle K_3 Y_l, Y_l\rangle = O(h^\infty)$. Consider the K_1 term. Writing $l = \alpha/h$ gives

$$\langle K_1 Y_l, Y_l\rangle = (2\pi h)^{-1}(2\pi)^{-1}\int_{\mathbb{R}}\int_0^{2\pi}\int_{\mathbb{R}} e^{i\big((\varphi-\varphi')v+G(v)-\alpha(\varphi-\varphi')\big)/h} a(\varphi-\varphi', v, h)\, dv\, d\varphi\, d\varphi'.$$

Changing integration variables to $(\varphi, \tilde\varphi = \varphi - \varphi')$, the kernel is independent of the first of these variables, so that integrating in it simply removes the factor 2π. We are left with

$$\langle K_1 Y_l, Y_l\rangle = (2\pi h)^{-1}\int_{\mathbb{R}}\int_{\mathbb{R}} e^{i(\tilde\varphi v + G(v) - \alpha\tilde\varphi)/h} a(\tilde\varphi, v, h)\, dv\, d\tilde\varphi.$$

The phase is stationary at the point $v = \alpha$, $-\tilde\varphi = G'(v) = \Sigma(v)$ and the stationary phase lemma shows that the integral is equal to

$$\langle K_1 Y_l, Y_l\rangle = e^{iG(\alpha)/h} a(-\Sigma(\alpha), \alpha, 0) + O(h) \tag{3.10}$$

(noting that the Hessian of the phase function has determinant 1 and signature 0).

Next we write

$$\langle K_2 Y_l, Y_l\rangle = (2\pi h)^{-1}(2\pi)^{-1}\int e^{i(\varphi-\varphi')v/h} b(\varphi, v, h) e^{-i\alpha(\varphi-\varphi')/h}\, dv\, d\varphi\, d\varphi'.$$

Here, the phase is stationary when $\alpha = v$. However, b is supported where $|v| \geq R_1 > R$ while $\alpha \leq R$ by hypothesis, so there are no stationary points on the support of the integrand. It follows that $\langle K_2 Y_l, Y_l\rangle = O(h^\infty)$. Thus by (3.9)

$$e^{i\beta_{l,h}} = e^{iG(\alpha)/h} a(-\Sigma(\alpha), \alpha, 0) + O(h). \tag{3.11}$$

The principal symbol of K_1 as an FIO is given as a multiple of the Liouville half-density on $\mathrm{T}^*\,\mathbb{S}^1$, $|d\varphi\, d\eta|^{1/2}$, by [10, Section 3]

$$\sigma(K_1)(\varphi, \eta, \varphi + \Sigma(\eta), -\eta) = a(-\Sigma(\eta), \eta, 0)|d\varphi\, d\eta|^{1/2}. \tag{3.12}$$

Indeed, the density d_C defined on page 143 of that paper equals $|d\varphi\, d\eta|$, where we used coordinates $(x, \theta) = (\varphi, \varphi', v)$. The principal symbol is the image of the map from C to Λ defined immediately following the definition of d_C, in the notation of that paper. In the notation of the current paper, $C = \mathrm{Crit}(\Phi)$ and Λ is the projection of the Legendrian L onto the first four coordinates. It follows from equation (3.12) and equation (3.6) that

$$a(-\Sigma(\eta), \eta, 0) = 1. \tag{3.13}$$

Combining (3.13) and (3.10), we see that

$$e^{i\beta_{l,h}} = e^{iG(\alpha)/h} + O(h), \tag{3.14}$$

establishing (3.2).

We proceed to the case $d = 3$. In this case, we will obtain the eigenvalue $e^{i\beta_{l,h}}$ by pairing the scattering matrix S_h with the highest weight spherical harmonics Y_l^l. These concentrate along a great circle γ, which we parametrize by arclength, $\varphi \in [0, 2\pi]$. Choose Euclidean coordinates in $\mathbb{R}^3$ so that the two-plane spanned by γ is the plane $x_3 = 0$. Then $Y_l^l = c_l(x_1 + ix_2)^l$ where c_l is a normalization factor, equal to $(2\pi)^{-1/2}(\pi l)^{1/4}(1 + O(l^{-1}))$. Let θ be the spherical coordinate equal to the angle with the positive x_3 axis. Then we can write

$$Y_l^l(\varphi, \theta) = c_l e^{il\varphi}(\sin\theta)^l = c_l e^{il\varphi} e^{-lg(\theta)} \tag{3.15}$$

where $g(\theta) = -\log \sin\theta = (\theta - \pi/2)^2/2 + O((\theta-\pi/2)^4)$.

In particular, expression (3.15) shows (and it is in any case well known) that the Y_l^l concentrate semiclassically (i.e. as $l \to \infty$) at the set $\{\theta = 0, \zeta = 0, \sigma = \alpha\}$ where $l = \alpha/h + O(1)$. Here we use coordinates (σ, ζ) dual to (φ, θ). So, to compute the pairing (3.9) with Y_l^l replacing Y_l, we first need to determine an oscillatory integral expression for K_1 that is valid in this microlocal region. (Note that the K_2 and K_3 terms give an $O(h^\infty)$ contribution as before.) So choose α_0 distance $\geq \epsilon$ from the set (3.3). As we will see, it suffices to find a local parametrization of L in a neighbourhood of

$$\{\theta = \theta' = \pi/2,\ \zeta = \zeta' = 0,\ \sigma = -\sigma' = \alpha_0,\ \varphi - \varphi' = \Sigma(\alpha_0)\};$$

this is the set of incoming and outgoing data of bicharacteristics which remain in the $x_3 = 0$ plane. To define this parametrization, we consider first a parametrization in two dimensions locally near a bicharacteristic where $\eta = \alpha$. As we have seen such a two dimensional parametrization is $(\varphi - \varphi')v + G(v)$, for v close to α. We note that when $v = \alpha$, $\varphi' - \varphi = \Sigma(\alpha)$, and we can write it in the form

$$\varphi' - \varphi = \pm \operatorname{dist}(\varphi, \varphi') + 2\pi k \tag{3.16}$$

for some integer k (recalling that the distance $\operatorname{dist}(\varphi, \varphi')$ lies strictly between 0 and π). We now claim that a suitable phase function is

$$\Phi(\omega, \omega', v) = (\mp \operatorname{dist}(\omega, \omega') - 2\pi k)v + G(v), \tag{3.17}$$

where $v \in \mathbb{R}$ is localized near α_0, $G(v)$ is as in (3.1), and the sign $\mp$ and the value of k agree with the two-dimensional case. Indeed, on each two-plane, if we use spherical coordinates $(\overline{\varphi}, \overline{\theta})$ adapted to that 2-plane then the form of the phase function agrees by construction with the two-dimensional phase function and therefore parametrizes that part of L associated to that 2-plane (since the dynamics on each 2-plane is identical to the $d = 2$ dynamics), that is, the subset (in the coordinates adapted to that 2-plane, indicated by a bar)

$$\{\overline{\theta} = \overline{\theta}' = \pi/2, \overline{\zeta} = \overline{\zeta}' = 0, \overline{\varphi}' - \overline{\varphi} = \Sigma(\alpha), \overline{\sigma} = -\overline{\sigma}' = \alpha, \tau = T(\alpha)\}. \tag{3.18}$$

We now observe that we can eliminate k by redefining $G(v)$ locally to be $G(v) + 2\pi k v$, which only has the irrelevant effect of changing Σ by $2\pi k$ (notice also that this does not affect the eigenvalue formula involving $e^{iG(lh)/h}$ in the statement of Proposition 3.1). From here on we only work with the $+$ sign in (3.16), i.e. the $-$ sign in (3.17), and $k = 0$. Notice that this means that $0 < \varphi' - \varphi < \pi$ and $0 < \Sigma(\alpha) < \pi$, i.e. $\sin \Sigma(\alpha) > 0$. Returning to our spherical coordinates associated to the 2-plane $x_3 = 0$, we can use the spherical cosine law applied to the spherical triangle with vertices (φ, θ), (φ', θ'), and the pole $x_3 = 1$:

$$\cos \operatorname{dist}((\varphi, \theta), (\varphi', \theta')) = (\cos(\varphi - \varphi') \sin\theta \sin\theta' + \cos\theta \cos\theta')$$

to write

$$\Phi(\varphi, y, \varphi', y, v) = -\cos^{-1} \left(\cos(\varphi - \varphi') \sin\theta \sin\theta' + \cos\theta \cos\theta'\right) v + G(v). \tag{3.19}$$

We can then write in these coordinates

$$\begin{aligned} L = \Big\{ &(\varphi, \theta, \varphi', \theta', \sigma, \zeta, \sigma', \zeta', \tau) \mid \\ &\sigma = \partial_\varphi \Phi = -\frac{v}{\sin \operatorname{dist}(\omega, \omega')} \left(\sin(\varphi - \varphi') \sin\theta \sin\theta' \right) \\ &\zeta = \partial_\theta \Phi = \frac{v}{\sin \operatorname{dist}(\omega, \omega')} \left(\cos(\varphi - \varphi') \cos\theta \sin\theta' - \sin\theta \cos\theta' \right) \\ &\sigma' = \partial_{\varphi'} \Phi = \frac{v}{\sin \operatorname{dist}(\omega, \omega')} \left(\sin(\varphi - \varphi') \sin\theta \sin\theta' \right) \\ &\zeta' = \partial_{\theta'} \Phi = \frac{v}{\sin \operatorname{dist}(\omega, \omega')} \left(\cos(\varphi - \varphi') \cos\theta' \sin\theta - \sin\theta' \cos\theta \right) \\ &\tau = \operatorname{dist}(\omega, \omega') v + G(v) \Big\} \quad \text{where } \operatorname{dist}(\omega, \omega') = G'(v). \end{aligned} \tag{3.20}$$

(Notice that by direct inspection we see that this agrees with (3.18) when $\theta = \theta' = \pi/2$, since then $\cos\theta = \cos\theta' = 0$ and $\operatorname{dist}(\omega, \omega') = \varphi' - \varphi = \Sigma(\alpha)$ and so $\sigma = v = \alpha$.)

The scattering matrix, microlocalized to this region of phase space, will then take the form

$$(2\pi h)^{-3/2} \int e^{i\Phi(\omega, \omega', v)/h} a(\omega, \omega', v, h) \, dv. \tag{3.21}$$

In terms of this parametrization the principal symbol of (3.21), say where both ω and ω' lie near the great circle γ and hence where we can use coordinates $(\varphi, \theta, \varphi', \theta'; \sigma, \zeta, \sigma', \zeta', \tau)$, is given at the point $(\varphi, \pi/2, \varphi + \Sigma(\alpha)h, \pi/2, \alpha, 0, -\alpha, 0, \tau(\alpha))$ by [10]

$$a(\varphi, \pi/2, \varphi + \Sigma(\alpha), \pi/2, \alpha, 0) e^{-i\pi/4} \big| ds \, d\theta \, d\sigma \, d\zeta \big|^{1/2} \Big| \det \frac{\partial(\varphi, \theta, \sigma, \zeta, d_v\Phi)}{\partial(\varphi, \theta, \varphi', \theta', v)} \Big|^{-1/2} \tag{3.22}$$

where the $e^{-i\pi/4}$ is a Maslov factor; see Remark 3.3 for more discussion about this. We need to compute the determinant above. We can disregard the repeated coordinates (φ, θ) and compute, using (3.20),

$$\det \frac{\partial(\sigma, \zeta, d_v\Phi)}{\partial(\varphi', \theta', v)} = \det \begin{pmatrix} 0 & 0 & -1 \\ 0 & \frac{-v}{\sin(\varphi' - \varphi)} & 0 \\ -1 & 0 & G''(v) \end{pmatrix} = \frac{v}{\sin(\varphi' - \varphi)} = \frac{\alpha}{\sin \Sigma(\alpha)} \text{ at } \theta = \theta' = \frac{\pi}{2}. \tag{3.23}$$

It follows that the principal symbol is

$$a(\varphi, \pi/2, \varphi + \Sigma(\alpha), \pi/2, \alpha, 0) e^{-i\pi/4} \Big(\frac{\alpha}{\sin \Sigma(\alpha)} \Big)^{-1/2} \big| ds \, d\theta \, d\sigma \, d\zeta \big|^{1/2}. \tag{3.24}$$

Then by equation (3.6)

$$a(\varphi, \pi/2, \varphi + \Sigma(\alpha), \pi/2, \alpha, 0) e^{-i\pi/4} = \Big(\frac{\alpha}{\sin \Sigma(\alpha)} \Big)^{1/2}. \tag{3.25}$$

We next write the contribution of K_1 to the expression (3.9) for the eigenvalue $e^{i\beta_{l,h}}$. Writing $l = \alpha/h$ and using (3.15) we get

$$\begin{aligned} \langle K_1 Y_l^l, Y_l^l \rangle = (2\pi h)^{-3/2} \int & e^{i\Phi(\varphi, \theta, \varphi', \theta', v)/h} e^{-i\alpha(\varphi - \varphi')/h} \left(\frac{\alpha}{\pi h} \right)^{1/2} (2\pi)^{-1} \\ & \times e^{-\alpha g(\theta)/h} e^{-\alpha g(\theta')/h} a(\varphi, \theta, \varphi', \theta', v, h) \, ds \, d\varphi' \, d\theta \, d\theta' \, dv \, (1 + O(h)) \end{aligned} \tag{3.26}$$

Here the factors $(\alpha/\pi h)^{1/2}(2\pi)^{-1}$ are to normalize the functions Y_l^l in L^2. We will analyze this using the stationary phase lemma with complex phase function, see e.g. [11, thm. 7.7.5]. Here the phase is

$$\Psi(\varphi,\theta,\varphi',\theta',v) = \Phi - \alpha(\varphi-\varphi') + i\alpha(g(\theta)+g(\theta')). \tag{3.27}$$

Notice that the integral as a function of (φ,φ') depends only on $\varphi-\varphi'$ by the rotational invariance of the scattering matrix, and the form of the Y_l^l which take the form $e^{il\varphi}$ times a function of θ. We change variable to $(\varphi,\tilde{\varphi})$, $\tilde{\varphi}=\varphi-\varphi'$ and integrate out the variable φ, giving us a factor of 2π. Then Ψ has nondegenerate stationary points in the remaining variables $(\tilde{\varphi},\theta,\theta',v)$. The imaginary part of the phase is stationary only at $\theta=\theta'=\pi/2$, while stationarity of the real part requires that $v=\alpha$ and $-\tilde{\varphi}=G'(v)=\Sigma(\alpha)$. The stationary phase lemma then gives us that (3.26) is equal to

$$\begin{aligned} &2\pi\left((2\pi h)^{-3/2}\left(\frac{\alpha}{\pi h}\right)^{1/2}(2\pi)^{-1}\right)(2\pi h)^2 \\ &\quad\times\left(e^{iG(\alpha)/h}\frac{1}{\det(-iD^2\Psi)^{1/2}}a(\varphi,\pi/2,\varphi+\Sigma(\alpha),\pi/2,\alpha,0)+O(h)\right). \end{aligned} \tag{3.28}$$

Here, to keep track of constants, we have written out all constants in (3.26); the first 2π comes from the integral in φ and the $(2\pi h)^2$ comes from the leading term in stationary phase in the four variables $(\tilde{\varphi},\theta,\theta',v)$. Simplifying the constants and using (3.25) this is equal to

$$\left(\frac{2\alpha^2}{\sin\Sigma(\alpha)}\right)^{1/2}\left(\frac{e^{iG(\alpha)/h}}{(\det -iD^2\Psi)^{1/2}}+O(h)\right). \tag{3.29}$$

We will show that, in the above expression

$$\det -iD^2\Psi(\varphi,0,\varphi+\Sigma(\alpha),0,\alpha) = \frac{2i\alpha^2}{\sin\Sigma(\alpha)}e^{-i\Sigma(\alpha)}. \tag{3.30}$$

Accepting this for the moment, we obtain from (3.28)

$$e^{i\beta_{l,h}} = e^{iG(\alpha)/h}e^{i\Sigma(\alpha)/2}+O(h).$$

Since $\Sigma(\alpha)=G'(\alpha)$ and $\alpha=lh$, this can be written

$$e^{i\beta_{l,h}} = e^{iG((l+1/2)h)/h}+O(h) \tag{3.31}$$

completing the proof of Proposition 3.1.

It remains to prove the formula for the Hessian in (3.30). First we notice that when $\theta=\theta'=\pi/2$ we have, using the formula in (3.19) for the distance function, in the coordinates $(\tilde{\varphi},\theta,\theta')$,

$$D^2\operatorname{dist}(\varphi,\pi/2,\varphi+\Sigma(\alpha),\pi/2) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \cot\Sigma(\alpha) & -\csc\Sigma(\alpha) \\ 0 & -\csc\Sigma(\alpha) & \cot\Sigma(\alpha) \end{pmatrix} \tag{3.32}$$

From this (3.17) and (3.27), we conclude that in the $(v,\tilde{\varphi},y,y')$ coordinates

$$D^2\Psi = \begin{pmatrix} G''(v) & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & -\alpha\cot\Sigma(\alpha)+i\alpha & \alpha\csc\Sigma(\alpha) \\ 0 & 0 & \alpha\csc\Sigma(\alpha) & -\alpha\cot\Sigma(\alpha)+i\alpha \end{pmatrix}. \tag{3.33}$$

Thus

$$\det -iD^2\Psi = -\alpha^2\Big(\cot^2\Sigma(\alpha) - 2i\cot\Sigma(\alpha) - 1 - \csc^2\Sigma(\alpha)\Big) = \frac{2i\alpha^2}{\sin\Sigma(\alpha)}e^{-i\Sigma(\alpha)}$$

and (3.30) holds. □

Remark 3.3. The Maslov factor in (3.22) and (3.25) arises as follows. First, Lemma 3.2 shows that the Maslov bundle over L is canonically trivial. However, unlike in the case $d = 2$, there is a nontrivial Maslov factor from comparing our phase function Φ above to one — let us call it $\tilde{\Phi}$ — that agrees with the canonical phase function, i.e. the pseudodifferential phase function, for $|\eta| \geq R$. By [10, Theorem 3.2.1], the principal symbol written relative to Φ contains the Maslov factor $e^{i\pi\sigma/4}$ where σ is the difference of signatures,

$$\sigma = \operatorname{sgn} D^2_{vv}\Phi - \operatorname{sgn} D^2_{\tilde{w}\tilde{w}}\tilde{\Phi}$$

where $\tilde{w} = (\tilde{w}_1, \tilde{w}_2)$ are the phase variables for $\tilde{\Phi}$. A tedious computation shows that $\sigma = -1$, leading to the Maslov factor in (3.22) and (3.25). (We remark that since Φ depends on one phase variable and $\tilde{\Phi}$ on two phase variables, by [10, Equation (3.2.12)] σ is odd, so the Maslov factor cannot vanish in this case.) Of course, the Maslov factors are irrelevant to the question of equidistribution, but they are relevant to the question of determining the eigenvalues modulo $O(h)$.

Proof of Proposition 1.3. In view of the remarks in the proof of Proposition 3.1, specifically equation (3.7), it is only necessary to do this in the cases $d = 2$ and $d = 3$. For definiteness, we write down the proof for $d = 3$; it is similar, and in fact simpler, for $d = 2$. Consider a spherical harmonic Y_l^l with $hl \geq R + h^\kappa$, where $\kappa < 1$. The eigenvalue $e^{i\beta_{l,h}}$ is given by (3.9) with Y_l^l replacing Y_l.

First assume that $hl \geq R' > R$. Then the K_1 term in (3.9) will be $O(h^\infty)$ (for a suitable decomposition of $S_h = K_1 + K_2 + K_3$ as above, with $R_2 < R'$), so we only have to consider the K_2 term. This is given by a pseudodifferential operator with symbol equal to $1 + O(h^\infty)$, so the $\langle Y_l^l, K_2 Y_l^l\rangle$ term is equal to $1 + O(h^\infty)$, proving the Proposition in this case.

Next assume that $R + h^\kappa \leq hl \leq R_1'$. For $R' < R_1$, the K_2 term in (3.9) will be $O(h^\infty)$ (for some other decomposition of S_h, with $R_1 > R'$), so we only need to consider the K_1 term. That is, it remains to show that

$$\langle (K_1 - \mathrm{Id})Y_l^l, Y_l^l\rangle = O(h^\infty) \text{ for } R + h^\kappa \leq hl$$

Using as above polar coordinates (φ, θ) on S^2 with dual coordinate (σ, ζ), we find a phase function Ψ for K_1 that parametrizes L microlocally in the region

$$\{|(\sigma,\zeta)|_g \geq R - \delta\}$$

for fixed small $\delta > 0$. Indeed, since L is given by the diagonal relation

$$\{\varphi = \varphi',\, \theta = \theta',\, \sigma = -\sigma',\, \zeta = -\zeta',\, \tau = 0\} \text{ for } \{|(\sigma,\zeta)|_g \geq R\}, \tag{3.34}$$

it follows that the functions $(\varphi, \theta, \sigma', \zeta')$ furnish local coordinates on the Legendrian L for $\{|(\sigma,\zeta)|_g \geq R\}$ and therefore, by continuity, for $\{|(\sigma,\zeta)|_g \geq R - \delta\}$ for some small $\delta > 0$. It then follows from [12, Theorem 21.2.18] that L can be parametrized by a phase function of the form

$$-\varphi' v - \theta' w + H(\varphi, \theta, v, w).$$

Since K_1 is pseudodifferential for $|(\sigma',\zeta')|_g^2 = \zeta'^2 + (\sin\theta')^{-2}\sigma'^2 \geq R^2$ (i.e. L satisfies (3.34)), we have

$$v \geq R\sin\theta' \implies H = \varphi v + \theta w \text{ and } b = 1 + O(h^\infty)$$

Thus

$$\langle (K_1 - \mathrm{Id}) Y_l^l, Y_l^l \rangle = \int \left(e^{i\left(-\varphi' v - \theta' w + H(\varphi,\theta,v,w)\right)/h} b(\varphi', \theta', v, w, h) - e^{i\left((\varphi-\varphi')v + (\theta-\theta')w\right)/h} \right) \times c_l e^{i\alpha(g(\theta)+g(\theta'))/h} \frac{d\varphi\, d\theta\, d\varphi'\, d\theta'\, dv\, dw}{(2\pi h)^3} + O(h^\infty), \tag{3.35}$$

As above we have written $l = \alpha/h$; hence $\alpha > R + h^\kappa$.

We insert cutoff functions by writing

$$1 = \chi\left(\frac{v - R\sin\theta'}{h^\kappa}\right) + (1-\chi)\left(\frac{v - R\sin\theta'}{h^\kappa}\right)$$

where $\chi(t)$ is supported in $t \le 1/2$, equal to 1 for $t \le 1/4$. With the cutoff χ inserted, the phase function is nonstationary on the support of the integrand, since stationarity requires that $v = \alpha$. It follows that we can integrate by parts arbitrarily many times, using the fact that the differential operator

$$\frac{1}{v - \alpha} \frac{h}{i} \frac{\partial}{\partial \varphi'}$$

leaves both exponential factors invariant; doing this gains a factor of $h^{1-\kappa}$ each time since $\alpha - v \ge h^\kappa/2$ on the support of the integrand. Thus the χ term is $O(h^\infty)$.

With the cutoff $1-\chi$ inserted, we write the integral in (3.35) in the form

$$\int e^{i\left((\varphi-\varphi')v + (\theta-\theta')w\right)/h} \left(e^{i\left(H(\varphi,\theta,v,w) - \varphi v - \theta w\right)/h} b(\varphi', \theta', v, w, h) - 1 \right) \times c_l (1-\chi)\left(\frac{v - R\sin\theta'}{h^\kappa}\right) e^{i\alpha(g(\theta)+g(\theta'))/h} \frac{d\varphi\, d\theta\, d\varphi'\, d\theta'\, dv\, dw}{(2\pi h)^3}. \tag{3.36}$$

We claim that the factor

$$\left(e^{i\left(H(\varphi,\theta,v,w) - \varphi v - \theta w\right)/h} b(\varphi', \theta', v, w, h) - 1 \right) \times (1-\chi)\left(\frac{v - R\sin\theta'}{h^\kappa}\right)$$

is $O(h^\infty)$. In fact, the term in the large brackets is $O(h^\infty)$ for $v \ge R\sin\theta'$, while if $v \le R\sin\theta'$, then the $1-\chi$ term vanishes identically. It follows that the $1-\chi$ term is also $O(h^\infty)$, completing the proof of Proposition 1.3. $\square$

4. EQUIDISTRIBUTION

For clarity of exposition, we will assume in this section that the energy E in (1.1) satisfies

$$E = 1.$$

The lemmas and propositions in this section apply to the arbitrary $E > 0$ case by writing $\widetilde{V} = V/E$ and $\widetilde{h} = h/\sqrt{E}$.

If $\omega = \left\{ e^{2\pi i x_1}, \ldots, e^{2\pi i x_K} \right\}$ is any set of K points on $\mathbb{S}^1$, then the **discrepancy** $D(\omega)$ is defined by

$$D(\omega) := \sup_{0 \le \phi_0 < \phi_1 \le 2\pi} \left| \frac{N(\phi_0, \phi_1; \omega)}{K} - \frac{\phi_1 - \phi_0}{2\pi} \right|, \tag{4.1}$$

where $N(\phi_0, \phi_1; \omega)$ is the number of points in ω with argument in $[\phi_0, \phi_1]$ (modulo 2π), counted with multiplicity. We state the following lemma in slightly more generality then is necessary for semiclassical potentials so that we may apply it without significant modification to the case of scattering by the disk.

Lemma 4.1. *Let $G(\alpha)\colon [0,R) \longrightarrow \mathbb{R}$ be smooth and assume that*

$$\{\alpha : G''(\alpha) = 0\} \text{ is finite in } [0,R).$$

Consider the points x_{lk} on the unit circle

$$\mathcal{E}_h = \{x_{lk} := \exp\left(i(G(lh)/h) : 0 \leq lh < R, k = 1, \dots, p_d(l)\right\}, \tag{4.2}$$

included according to multiplicity. Here $p_d(l) = \dim\ker\left(\Delta_{\mathbb{S}^{d-1}} - l(l+d-2)\right)$.

Then the sets $\mathcal{E}_h$ equidistribute as $h \downarrow 0$. That is, the discrepancy satisfies

$$\lim_{h\downarrow 0} D(\mathcal{E}_h) = 0. \tag{4.3}$$

To apply the lemma to the eigenvalues of the scattering matrix S_h, we must show that they still equidistribute despite satisfying only the weaker asymptotic condition in Proposition 3.1. Below, S will denote any set satisfying

1. S is a finite subset of $[0,R]$.
2. If $0 \leq \alpha \leq R$ and $G''(\alpha) = 0$, then $\alpha \in S$.
3. $0, R \in S$.

For example, to apply Proposition 4.2 to Proposition 3.1, when the dimension d is odd one takes S to be the set to be

$$\{\alpha \leq R : G''(\alpha) = 0\} \cup \{\alpha \leq R : G'(\alpha) = 0\} \cup \{0, R\}.$$

Proposition 4.2. *Notation and assumptions as in Lemma 4.1, let*

$$\widetilde{\mathcal{E}}_h = \{\widetilde{x}_{lk} : 0 \leq lh \leq R, k = 1, \dots, p_d(l)\}, \tag{4.4}$$

be a collection of points on $\mathbb{S}^1$ (included according to multiplicity), such that for any $\epsilon > 0$, if l satisfies $\operatorname{dist}(lh, S) \geq \epsilon$ then

$$\widetilde{x}_{lk} = \exp i(G(lh)/h) + E(l,h))$$

where $|E(l,h)| < C(\epsilon)h$. Then, if G satisfies the condition in Lemma 4.1,

$$\lim_{h\downarrow 0} D(\widetilde{\mathcal{E}_h}) = 0.$$

We will use the following notation. With any set S as above, let

$$\begin{aligned} \mathcal{E}_h(\epsilon) &:= \mathcal{E}_h \cap \{x_{lk} \colon \operatorname{dist}(lh, S) \geq \epsilon\} \\ \widetilde{\mathcal{E}}_h(\epsilon) &:= \widetilde{\mathcal{E}}_h \cap \{\widetilde{x}_{lk} \colon \operatorname{dist}(lh, S) \geq \epsilon\}, \end{aligned} \tag{4.5}$$

always understood to include points according to multiplicity.

Proof of Proposition 4.2 assuming Lemma 4.1: By Proposition 3.1, for every $\epsilon > 0$, there is a constant $C = C(\epsilon, S) > 0$ so that for h sufficiently small,

$$N(\phi_0 + Ch, \phi_1 - Ch; \mathcal{E}_h(\epsilon)) \leq N(\phi_0, \phi_1; \widetilde{\mathcal{E}_h}(\epsilon)) \leq N(\phi_0 - Ch, \phi_1 + Ch; \mathcal{E}_h(\epsilon)).$$

More precisely, C depends on the number of points in $S \cap [\epsilon, R-\epsilon]$. Dividing through by $2(R/h)^{d-1}/(d-1)!$ and subtracting $(\phi_1 - \phi_0)/2\pi$ gives

$$\begin{aligned}\left| \frac{N(\phi_0, \phi_1; \widetilde{\mathcal{E}_h}(\epsilon))}{2(R/h)^{d-1}/(d-1)!} - \frac{\phi_1 - \phi_0}{2\pi} \right| &\leq \max \left\{ \left| \frac{N(\phi_0 + Ch, \phi_1 - Ch; \mathcal{E}_h(\epsilon))}{2(R/h)^{d-1}/(d-1)!} - \frac{\phi_1 - \phi_0}{2\pi} \right|, \right. \\ &\qquad \left. \left| \frac{N(\phi_0 - Ch, \phi_1 + Ch; \mathcal{E}_h(\epsilon))}{2(R/h)^{d-1}/(d-1)!} - \frac{\phi_1 - \phi_0}{2\pi} \right| \right\} \\ &\leq D(\mathcal{E}_h(\epsilon)) + (1 + C(\epsilon))O(h) + O(\epsilon),\end{aligned}$$

uniformly in h and ϵ, where for the second inequality we used

$$|\mathcal{E}_h(\epsilon)| = \frac{2R^{d-1}}{h^{d-1}(d-1)!} \left(1 + O(h) + O(\epsilon)\right), \tag{4.6}$$

where $|\mathcal{E}_h(\epsilon)|$ is the number of points in $\mathcal{E}_h(\epsilon)$. Similarly, $D(\mathcal{E}_h) = D(\mathcal{E}_h(\epsilon)) + O(h) + O(\epsilon)$ for h, ϵ small. The same is true for $\widetilde{\mathcal{E}}_h$. Thus

$$\left| \frac{N(\phi_0, \phi_1; \widetilde{\mathcal{E}_h})}{2(R/h)^{d-1}/(d-1)!} - \frac{\phi_1 - \phi_0}{2\pi} \right| \leq D(\mathcal{E}_h) + (1 + C(\epsilon))O(h) + O(\epsilon). \tag{4.7}$$

Thus

$$\limsup_{h \downarrow 0} D(\widetilde{\mathcal{E}}_h) = O(\epsilon),$$

and as $\epsilon > 0$ was arbitrary, we obtain the result. □

Remark 4.3. Note that the proof gives no information about the exact vanishing rate of $D(\widetilde{\mathcal{E}}_h)$ as $h \to 0$. For this, one must have information on the dependence of $C(\epsilon)$ on ϵ, and then optimize in ϵ in (4.7) as $h \to 0$. This is what we do in Section 6 to obtain improved remainders in the case of scattering by the disk.

To prove the lemma, we use theorems from [13]. The following theorem follows from [13, ch. 2, eq. 2.42]

Theorem 4.4 (Erdös-Turán). *There is a constant $c > 0$ such that if*

$$\omega = \left\{ e^{2\pi i x_1}, \ldots, e^{2\pi i x_N} \right\}$$

is a finite sequence of N points on $\mathbb{S}^1$ and m is any positive integer, then

$$D(\omega) \leq c \left(\frac{1}{m} + \sum_{j=1}^{m} \frac{1}{j} \left| \frac{1}{N} \sum_{l=1}^{N} e^{2\pi i j x_l} \right| \right). \tag{4.8}$$

To bound the exponential sums that appear on the right hand side of (4.8), we use [13, ch. 1, thm. 2.7], namely

Theorem 4.5. *Let a and b be integers with $a < b$, and let f be twice differentiable on $[a, b]$ with $|f''(x)| \geq \rho > 0$ for $x \in [a, b]$. Then*

$$\left| \sum_{l=a}^{b} e^{2\pi i f(l)} \right| \leq (|f'(b) - f'(a)| + 2) \left(\frac{4}{\sqrt{\rho}} + 3 \right). \tag{4.9}$$

Finally there is [13, thm. 2.6] (with minor modifications in notation)

Theorem 4.6. *For $1 \le i \le k$, let ω_i be a set of $|\omega_i|$ points on $\mathbb{S}^1$ with discrepancy $D(\omega_i)$. Let ω be a concatenation of $\omega_1, \dots, \omega_k$, that is, a set obtained by listing in some order the terms of the ω_i. Then*

$$D(\omega) \le \sum_{i=1}^{k} \frac{|\omega_i|}{|\omega|} D(\omega_i), \tag{4.10}$$

where $|\omega|$ is the number of points in ω.

Proof of Lemma 4.1. We begin by assuming that G'' has no zeroes in the open interval $(0, R)$. In this case, the set S in (4.5) above is equal to $\{0, R\}$.

We first analyze the subset $\mathcal{E}_h(\epsilon) \subset \mathcal{E}_h$ defined in (4.5). Define

$$\begin{aligned} \tilde\rho = \tilde\rho(\epsilon) &= \min_{\epsilon \le \alpha \le R-\epsilon} |G''(\alpha)| \\ \tilde\kappa = \tilde\kappa(\epsilon) &= 2 \max_{\epsilon \le \alpha \le R-\epsilon} |G'(\alpha)| . \end{aligned} \tag{4.11}$$

We will show that for each $\gamma \in (0,1)$ there is a constant $c = c(\gamma) > 0$ so that for each $\epsilon > 0$,

$$D(\mathcal{E}_h(\epsilon)) < c \left(h^\gamma + \tilde\kappa \tilde\rho^{-1/2} h^{1/2 - \gamma/2} + \tilde\rho^{-1/2} h^{1/2} + \tilde\kappa h^{1-\gamma} \right) \tag{4.12}$$

Since $\mathcal{E}_h(\epsilon) - \mathcal{E}_h = h^{-d+1}(O(\epsilon) + O(h))$, for some $c = c(\gamma) > 0$ independent of ϵ we have.

$$\begin{aligned} D(\mathcal{E}_h) = &\sup_{0 \le \phi_0 < \phi_1 \le 2\pi} \left| \frac{N(\phi_0, \phi_1; \mathcal{E}_h)}{|\mathcal{E}_h|} - \frac{\phi_1 - \phi_0}{2\pi} \right| \\ &\le c(\epsilon + h) + D(\mathcal{E}_h(\epsilon)) \end{aligned} \tag{4.13}$$

So

$$\limsup_{h \downarrow 0} D(\mathcal{E}_h) \le c\epsilon,$$

and as $\epsilon > 0$ was arbitrary, we obtain (4.3).

Case 1: dimension $d = 2$. Note that when $d = 2$ the multiplicity of the eigenspaces is $p_2(l) = 1$ if $l = 0$ and 2 otherwise, so that

$$|\mathcal{E}_h(\epsilon)| = 2 \left(\lfloor (R-\epsilon)/h \rfloor - \lceil \epsilon/h \rceil + 1 \right).$$

We apply Theorem 4.4 with $\omega = \mathcal{E}_h(\epsilon)$, so that, in the notation of Theorem 4.4, $x_l = G(lh)/(2\pi h)$. Thus

$$D(\mathcal{E}_h(\epsilon)) \le c \left(\frac{1}{m} + \sum_{j=1}^{m} \frac{1}{j} \left| \frac{1}{\lfloor (R-\epsilon)/h \rfloor - \lceil \epsilon/h \rceil + 1} \sum_{l=\lceil \epsilon/h \rceil}^{\lfloor (R-\epsilon)/h \rfloor} e^{ijG(lh)/h} \right| \right)$$

Then we apply Theorem 4.5 with $f(x) = (j/2\pi) G(xh)/h$, $a = \lceil \epsilon/h \rceil$, and $b = \lfloor (R-\epsilon)/h \rfloor$. Thus, if $xh \le R - \epsilon$ then $|f''(x)| = hj\,|G''(xh)|\,/2\pi \ge hj\tilde\rho/2\pi$, which equals ρ in the notation of Theorem 4.5. It follows that

$$D(\mathcal{E}_h(\epsilon)) \le c \left(\frac{1}{m} + \frac{h}{R - 2\epsilon - h} \sum_{j=1}^{m} \frac{1}{j} \left(\frac{j\tilde\kappa}{\pi} + 2 \right) \left(\left(\frac{32\pi}{hj\tilde\rho} \right)^{1/2} + 3 \right) \right). \tag{4.14}$$

By letting $m = \lfloor h^{-\gamma} \rfloor$ for any $\gamma > 0$, we obtain (4.12).

Finally, suppose there are a finite number of points $0 < a_1 < \dots < a_{n-1} < R$ with $G(\alpha_i = 0$, and let $a_0 = 0, a_n = R$. Note that, if we define $\mathcal{E}_h(a, b)$ to be the set of $x_{l,k}$ with $a \le lh \le b$, counted with multiplicity. Then the above arguments show that

$\lim_h D(\mathcal{E}_h(a,b)) = 0$, in fact if $\widetilde{\rho}(\epsilon, h)$ (resp. $\widetilde{\kappa}$) is defined to be $\min_{a+\epsilon\leq\alpha\leq b-\epsilon} |G''(\alpha)|$ (resp. min), then the proof is the same. The lemma in the $d = 2$ case now follows from Theorem 4.6 since by this equation

$$D(\mathcal{E}_h) \leq \sum_{i=1}^{n} D(\mathcal{E}_h(a_{i-1}, a_i)).$$

The proof is now complete in the case $d = 2$.

Case 2: dimension $d > 2$. As in the $d = 2$ case, we begin by assuming that $G''(\alpha)$ has zeroes only at 0 and R.

We will apply Theorem 4.6 to $D(\mathcal{E}_h(\epsilon))$ decomposed as a superposition in the following way. It will be convenient to set

$$N := \lfloor (R-\epsilon)/h \rfloor. \tag{4.15}$$

Define,

$$\omega(n) := \left\{ e^{iG(lh)/h} : n \leq l \leq N \rfloor \right\}$$

with *unit multiplicity*. Note that $\omega(n)$ has $N - n + 1$ elements. Setting

$$\begin{aligned}
\omega_1 &= \omega(0)\\
\omega_2 &= \omega(1)\\
&\vdots\\
\omega_{p_d(1)} &= \omega(1)\\
\omega_{p_d(1)+1} &= \omega(2)\\
&\vdots\\
\omega_{p_d(2)} &= \omega(2)\\
&\vdots\\
\omega_{p_d(N-1)+1} &= \omega(N)\\
&\vdots\\
\omega_{p_d(N)} &= \omega(N),
\end{aligned}$$

we see that the set $\mathcal{E}_h(\epsilon)$ is the superposition of the sets $\omega_1, \ldots \omega_{p_d(N)}$.

The discrepency $D(\omega(n))$ can be estimated using the method from the $d = 2$ case. In particular, as in (4.14) we see that for any positive integer m,

$$D(\omega(n)) \leq c \left(\frac{1}{m} + \sum_{j=1}^{m} \frac{1}{(N-n+1)j} \left| (\tilde{\kappa} j + 2) \left(\left(\frac{32}{jc\tilde{\rho}h} \right)^{1/2} + 3 \right) \right| \right) \tag{4.16}$$

By Theorem 4.6, we have

$$\begin{aligned}D(\mathcal{E}_h(\epsilon)) &\leq \sum_{i=0}^{p_d(N)} \frac{|\omega_i|}{|\mathcal{E}_h(\epsilon)|} D(\omega_i)\\ &\leq \frac{1}{|\mathcal{E}_h(\epsilon)|} \sum_{n=0}^{N} |\omega(n)| \left(p_d(n) - p_d(n-1)\right) D(\omega(n))\\ &\leq ch^{d-1} \sum_{n=1}^{N} (N-n+1)(n+1)^{d-3} D(\omega(n)),\end{aligned} \tag{4.17}$$

Substituting the estimate (4.16) into (4.17), again with $m = \lfloor h^{-\gamma} \rfloor$ for some fixed $\gamma \in (0,1)$, we end up with five terms to deal with corresponding to the five terms in the right hand side of (4.16). For all of these we use standard bounds for sums of polynomials and $N \sim c/h$. The easiest is the $1/m$ term, since

$$h^{d-1} \sum_{n=1}^{N} (N-n+1)(n+1)^{d-3} h^{\gamma} \leq ch^{\gamma}. \tag{4.18}$$

Next we do the terms involving $\tilde{\rho}$. There is

$$\begin{aligned}&h^{d-1} \sum_{n=1}^{N} (N-n+1)(n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} \frac{1}{(N-n+1)j} (\tilde{\kappa} j) \left(\frac{32}{jc\tilde{\rho}h}\right)^{1/2}\\ &\leq ch^{d-1} \tilde{\kappa} \sum_{n=1}^{N} (n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} \left(\frac{32}{j\tilde{\rho}h}\right)^{1/2}\\ &\leq ch^{d-1} \tilde{\kappa} \left(\frac{1}{\tilde{\rho}h}\right)^{1/2} \sum_{n=1}^{N} (n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} j^{-1/2}\\ &\leq c \left(\frac{1}{\tilde{\rho}}\right)^{1/2} \tilde{\kappa} h^{1/2-\gamma/2},\end{aligned} \tag{4.19}$$

and

$$\begin{aligned}&h^{d-1} \sum_{n=1}^{N} (N-n+1)(n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} \frac{2}{(N-n+1)j} \left(\frac{32}{jc\tilde{\rho}h}\right)^{1/2}\\ &\leq h^{d-1} c \left(\frac{1}{\tilde{\rho}h}\right)^{1/2} \sum_{n=1}^{N} (n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} j^{-3/2}\\ &\leq c \left(\frac{1}{\tilde{\rho}}\right)^{1/2} h^{-1/2}.\end{aligned} \tag{4.20}$$

The other terms are

$$\begin{aligned}&h^{d-1} \sum_{n=1}^{N} (N-n+1)(n+1)^{d-3} \sum_{j=1}^{\lfloor h^{-\gamma} \rfloor} \frac{1}{(N-n+1)j} (\tilde{\kappa} j) \times 3\\ &\leq c\tilde{\kappa} h^{1-\gamma} \leq c\tilde{\kappa} h^{1/2-\gamma/2},\end{aligned} \tag{4.21}$$

and

$$h^{d-1}\sum_{n=1}^{N}(N-n+1)(n+1)^{d-3}\sum_{j=1}^{\lfloor h^{-\gamma}\rfloor}\frac{6}{(N-n+1)j}\leq ch\log(1/h), \tag{4.22}$$

Combining (4.18)– (4.22) with (4.17) gives (4.12).

We take care of the case of a non-trivial number of zeroes of G'' on $[0,R]$ exactly as in the $d=2$ case.

□

We can now prove Theorem 1.1.

Proof of Theorem 1.1. Theorem 1.1 is an direct consequence of Proposition 3.1, Lemma 4.1, and Proposition 4.2. □

5. Examples of potentials that satisfy assumption 1.8

We begin by deriving an integral expression for the scattering angle, $\Sigma(\alpha)$. Note that, if the potential is non-trapping at energy E, then along a bicharacteristic, the functions ρ and $\dot{\rho}$ do not any have simulateous zeros. For if there were such a time, and the value of r at this time were r_0, then $r(t)\equiv r_0$ would be a bicharacteristic, contradicting the non-trapping assumption. Hence the zeros of the function $E-\alpha^2/r^2-V(r)$ are simple on the region of interaction (1.4). Given a fixed bicharacteristic, let r_m be the minimum value of r; note that r_m is a function only of the angular momentum α. We denote the derivative of r_m with respect to α by $r_m'(\alpha)$. By symmetry, this is the unique point at which $\rho=0$ and $\dot{\rho}>0$, so we can divide the bicharacteristic 'in half,' and consider only times when $r>r_m$ and $\rho\geq 0$. We write

$$\frac{d\varphi}{dr}=\frac{\alpha/r^2}{\rho}=\frac{\alpha}{r^2\sqrt{E-\alpha^2/r^2-V(r)}}.$$

By the simplicity of the zeros in the denominator, we can integrate to obtain, for $\alpha>0$,

$$\Sigma(\alpha)=\pi-2\int_{r_m}^{\infty}\frac{\alpha}{r^2\sqrt{E-\alpha^2/r^2-V(r)}}dr. \tag{5.1}$$

(See Figure 1. The change in θ is the angle between the incoming and outgoing trajectories.)

Proposition 5.1. *Suppose that on the region of interaction $\mathcal{R}$ the potential V satisfies, $V'\leq 0$ (so necessarily $V\geq 0$ since V is compactly supported) and $r(V')^2+(E-V)(V'+rV'')\geq 0$ with equality only when $V=0$. Then $\partial_\alpha\Sigma(\alpha)<0$ for for $\alpha\in[0,R)$.*

The same is true if $V=cW$, where c is sufficiently large and where $W(r)=0$ for $r\geq R$, $W(r)>0$ for $0\leq r<R$ and $W''(r)$ is positive and monotone decreasing in some nonempty interval $[R-\epsilon,R)$.

In (5.1), set $s=r/r_m$, so

$$\begin{aligned}\Sigma(\alpha)&=\pi-2\int_{r_m}^{\infty}\frac{\alpha}{(sr_m)^2\sqrt{E-\alpha^2/((sr_m)^2-V(sr_m)}}d(sr_m)\\&=\pi-2\int_{1}^{\infty}\frac{\alpha}{s^2\sqrt{Er_m^2-\alpha^2/s^2-r_m^2V(sr_m)}}ds.\end{aligned}$$

Differentiating under the integral sign gives

$$\begin{aligned} -\frac{1}{2}\Sigma'(\alpha) &= \int_1^\infty \Bigg(\frac{1}{s^2\sqrt{Er_m^2 - \alpha^2/s^2 - r_m^2 V(sr_m)}} \\ &\qquad -\frac{\alpha}{2}\,\frac{2r_m r'_m - 2\alpha/s^2 - 2r_m V(sr_m) r'_m - r_m^2 s V'(sr_m) r'_m}{s^2\left(Er_m^2 - \alpha^2/s^2 - r_m^2 V(sr_m)\right)^{3/2}} \Bigg) ds \\ &= \int_1^\infty \frac{\big(r_m^2(E - V(sr_m)\big) - \alpha r_m r'_m(E - V(sr_m) - \frac{1}{2} r_m s V'(sr_m))}{s^2\left(Er_m^2 - \alpha^2/s^2 - r_m^2 V(sr_m)\right)^{3/2}}\, ds \end{aligned}$$

Differentiating

$$E - \alpha^2/r_m^2 - V(r_m) = 0 \tag{5.2}$$

shows $\alpha r_m r'_m = \frac{\alpha^2}{E - V(r_m) - \frac{1}{2} r_m V'(r_m))}$. Plugging this in gives

$$\begin{aligned} -\frac{1}{2}\Sigma'(\alpha) = \int_1^\infty & \left(r_m^2 (E - V(sr_m)) - \alpha^2 \frac{E - V(sr_m) - \frac{1}{2} r_m s V'(sr_m)}{E - V(r_m) - \frac{1}{2} r_m V'(r_m)} \right) \\ & \times \frac{1}{s^2\left(Er_m^2 - \alpha^2/s^2 - r_m^2 V(sr_m)\right)^{3/2}}\, ds \end{aligned}$$

and using (5.2) again shows that $(1/2)\partial_\alpha \Sigma(\alpha)$ is equal to

$$\begin{aligned} & \int_1^\infty \frac{E - V(sr_m) - \frac{1}{2} r_m s V'(sr_m)}{s^2\left(Er_m^2 - \alpha^2/s^2 - r_m^2 V(sr_m)\right)^{3/2}} \\ & \quad \times \left(\frac{E - V(sr_m)}{E - V(sr_m) - \frac{1}{2} r_m s V'(sr_m)} - \frac{E - V(r_m)}{E - V(r_m) - \frac{1}{2} r_m V'(r_m)} \right) r_m^2 ds \\ & = \int_{r_m}^\infty \frac{E - V(r) - \frac{1}{2} r V'(r)}{r^2\left(E - \alpha^2/r^2 - V(r)\right)^{3/2}} \\ & \quad \times \left(\frac{E - V(r)}{E - V(r) - \frac{1}{2} r V'(r)} - \frac{E - V(r_m)}{E - V(r_m) - \frac{1}{2} r_m V'(r_m)} \right) dr. \end{aligned}$$

Differentiating the expression $(E - V(r))/(E - V(r) - \frac{1}{2} r V'(r))$ with respect to r and using $V' \leq 0$, we see that the integrand is positive if for $r_m < r < R$ if

$$r(V')^2 + (E - V)(V' + rV'') > 0. \tag{5.3}$$

This is exactly the condition in the first paragraph of the proposition.

To check that the condition in the second paragraph is also sufficient, observe the following. If $V(r) \geq 0$, $V'(r) < 0$ and $V''(r) > 0$ on some open interval $(R - \epsilon, R)$, it follows that for r sufficiently close to R, that $V' + rV'' > 0$. By picking large enough $c > 0$, (5.3) will hold on the region of interaction $\mathcal{R}$.

6. Scattering by the disk

In this section we will prove Theorem 1.5 from the introduction. We restrict our attention to the disk of radius 1, since the phase shifts for the disk of radius R can be obtained from those for $R = 1$ by a scalling argument.

Here we use on an ODE analogous to that in (1.3) to give a formula for the eigenvalues. In fact, for any smooth solution f_l to $\Delta_{\mathbb{S}^{d-1}} f_l = l(l+d-2) f_l$, a straightforward computation shows that

$$S_k(f_l) = -\frac{H^{(1)}_{l+(d-2)/2}(k)}{H^{(2)}_{l+(d-2)/2}(k)} f_l, \tag{6.1}$$

where the $H^{(i)}_\nu$ are Hankel functions of order ν [1].

For $z > 0$ define $\zeta = \zeta(z)$ by

$$\begin{aligned} \frac{2}{3}(\zeta)^{3/2} &= \int_z^1 \frac{\sqrt{1-t^2}}{t}dt = \log\frac{1+\sqrt{1-z^2}}{z} - \sqrt{1-z^2}, \\ \frac{2}{3}(-\zeta)^{3/2} &= \int_1^z \frac{\sqrt{t^2-1}}{t}dt = \sqrt{z^2-1} - \cos^{-1}\left(\frac{1}{z}\right). \end{aligned} \tag{6.2}$$

The two formulas for ζ are equivalent; the first is more convenient for $z < 1$ and the second for $z > 1$.

Lemma 6.1. *As* $k, l \to \infty$

$$\arg H^{(1)}_\nu(k) = \begin{cases} \frac{-\pi}{2} + O((\nu^{-1}\zeta^{-3/2})^\infty) + O(\nu^{-\infty}), & k/\nu < 1, \\ \frac{-5\pi}{12} + \arg\left(\exp\left(-\frac{2}{3}\nu\zeta^{3/2}\right)\right) + O(\nu^{-1}\zeta^{-3/2}) + O(\nu^{-1}\zeta^{1/2}) & \\ \qquad +O(\nu^{-\infty}), & k/\nu > 1 \end{cases} \tag{6.3}$$

where $\nu = l + (d-2)/2$, *and*

$$\zeta = \zeta(k/\nu)$$

(see (6.2)*).*

In particular, for any $p \in (1/3, 1)$, *if we restrict to the regions* $k^p < \nu < k - k^p$ *or* $\nu > k + k^p$, *then the error terms in* (6.3) *becomes*

$$\arg H^{(1)}_\nu(k) = \begin{cases} \frac{-\pi}{2} + O(k^{-\infty}), & \nu > k + k^p, \\ \frac{-5\pi}{12} + \arg\left(\exp\left(-\frac{2}{3}\nu\zeta^{3/2}\right)\right) + O(k^{(1-3p)/2}), & k^p < \nu < k - k^p \end{cases} \tag{6.4}$$

We will prove this in a moment. Using Lemma 6.1 and the arguments above, we can now prove Theorem 1.5.

Proof of Theorem 1.5: First we will bound $D(\mathcal{E}_{1/k}(\epsilon))$, where $\mathcal{E}_{1/k}(\epsilon)$ is as in (4.2) $G(\alpha) = -\frac{2}{3}\alpha\zeta(1/\alpha)^{3/2}$ and $\alpha = \nu/k$. We apply (4.12), using

$$\begin{aligned} \widetilde{\kappa} &\le \max_{0\le\alpha\le1} G' \le c < \infty \\ \widetilde{\rho} &\ge \min_{0\le\alpha\le1} G''(\alpha) \ge c > 0. \end{aligned} \tag{6.5}$$

This means that in (4.12), $\tilde{\rho}(\epsilon, 1/k) \ge c > 0$ and $\tilde{\kappa}(\epsilon, 1/k) \le c < \infty$ for all ϵ, and thus for any $0 < \gamma < 1$,

$$D(\mathcal{E}_{1/k}(\epsilon)) \le c\left(k^{-\gamma} + k^{-1/2+\gamma/2}\right). \tag{6.6}$$

Now we bound the true driscrepancy $\widetilde{\mathcal{E}}_{1/k}$ by following the proof of Proposition 4.2 with $k = 1/h$, specifically using (4.7) and $C(\epsilon)h = O(k^{(1-3p)/2})$, $\epsilon = k^{p-1}$ and taking $\gamma = 1/3$ in (6.6) gives (1.12).

□

Proof of Lemma 6.1: We will use the following uniform asymptotic (9.3.37 from [1]). For $\zeta = \zeta(z)$ as in (6.2),

$$\begin{aligned} H^1_\nu(\nu z) = 2e^{-\pi i/3}\left(\frac{4\zeta}{1-z^2}\right)^{1/4}&\left(\frac{\operatorname{Ai}\left(e^{2\pi i/3}\nu^{2/3}\zeta\right)}{\nu^{1/3}} + \frac{e^{2\pi i/3}\operatorname{Ai}'\left(e^{2\pi i/3}\nu^{2/3}\zeta\right)}{\nu^{5/3}} b_0(\zeta)\right) \\ &\times\left(1 + O_R(\nu^{-2}) + O(\nu^{-\infty})\right), \end{aligned} \tag{6.7}$$

where $\nu \to \infty$, $z \in (0,\infty)$, b_0 is bounded and positive, and O_R denotes a real-valued remainder.

The Airy function $\operatorname{Ai}(x)$ satisfies

$$\begin{aligned} \operatorname{Ai}(x) &\sim \frac{1}{2}\pi^{-1/2}x^{-1/4}e^{-2x^{3/2}/3}\sum_{k=0}^{\infty} c_k x^{-3k/2} \\ \operatorname{Ai}'(x) &\sim -\frac{1}{2}\pi^{-1/2}x^{1/4}e^{-2x^{3/2}/3}\sum_{k=0}^{\infty} d_k x^{-3k/2}, \end{aligned} \tag{6.8}$$

as $|x| \to \infty$ with $|\arg x| < \pi$ (chapter 10 of [1]). The c_k and d_k are real, and $c_0 = d_0 = 1$.

We will plug (6.8) into (6.7) with $x = e^{2\pi i/3}\nu^{2/3}\zeta$.

$$\begin{aligned} \arg H^1_\nu(\nu z) &= -\frac{\pi}{3} + \arg\left(\frac{\operatorname{Ai}\left(e^{2\pi i/3}\nu^{2/3}\zeta\right)}{\nu^{1/3}} + \frac{e^{2\pi i/3}\operatorname{Ai}'\left(e^{2\pi i/3}\nu^{2/3}\zeta\right)}{\nu^{5/3}} b_0(\zeta)\right) + O(\nu^{-\infty}) \\ &= -\frac{\pi}{3} + \arg\operatorname{Ai}(x) + O(\nu^{-\infty}) + \arg\left(1 - \left(\frac{e^{2\pi i/3}\operatorname{Ai}'(x)}{\operatorname{Ai}(x)\nu^{4/3}}\right)\right) + O(x^{-\infty}). \end{aligned} \tag{6.9}$$

Consider first the case $z < 1$. Then $\zeta > 0$. Using that $\arg\operatorname{Ai}(x) = -\pi/6 + O(x^{-\infty})$ and that $\operatorname{Im}(e^{2\pi i/3}\operatorname{Ai}'(x)/\operatorname{Ai}(x)) = O(x^{-\infty})$, we conclude that

$$\arg H^1_\nu(\nu z) = -\pi/2 + O(\nu^{-\infty}) + O(x^{-\infty}). \tag{6.10}$$

Now consider the case $z > 1$. In (6.9) we take an additional step

$$\begin{aligned} \arg H^1_\nu(\nu z) &= -\frac{\pi}{3} + \arg\operatorname{Ai}(x) + O(\nu^{-\infty}) + \arg\left(1 - \left(\frac{e^{2\pi i/3}\operatorname{Ai}'(x)}{\operatorname{Ai}(x)\nu^{4/3}}\right)\right) + O(x^{-\infty}). \\ &= -\frac{\pi}{3} + \arg\left(x^{1/4}e^{-2\nu\zeta^{3/2}/3}\left(1 + O(x^{-3/2})\right)\right) \\ &\quad + O(\nu^{-\infty}) + \arg\left(1 - \left(\frac{e^{2\pi i/3}\operatorname{Ai}'(x)}{\operatorname{Ai}(x)\nu^{4/3}}\right)\right). \\ &= -\frac{\pi}{3} + \arg\left(x^{1/4}e^{-2\nu\zeta^{3/2}/3}\right) + O(x^{-3/2}) \\ &\quad + O(\nu^{-\infty}) + O(x^{1/2}\nu^{-4/3}). \end{aligned} \tag{6.11}$$

This completes the proof of (6.3).

□

References

[1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.

[2] I. Alexandrova. Structure of the semi-classical amplitude for general scattering relations. *Comm. Partial Differential Equations*, 30(10-12):1505–1535, 2005.

[3] M. S. Birman and D. R. Yafaev. Asymptotics of the spectrum of the s-matrix in potential scattering. *Soviet Phys. Dokl.*, 25(12):989–990, 1980.
[4] M. S. Birman and D. R. Yafaev. Asymptotic behavior of limit phases for scattering by potentials without spherical symmetry. *Theoret. Math. Phys.*, 51(1):344–350, 1982.
[5] M. S. Birman and D. R. Yafaev. Asymptotic behaviour of the spectrum of the scattering matrix. *J. Sov. Math.*, 25:793–814, 1984.
[6] M. S. Birman and D. R. Yafaev. Spectral properties of the scattering matrix. *St Petersburg Math. Journal*, 4(6):1055–1079, 1993.
[7] D. Bulger and A. Pushnitski. The spectral density of the scattering matrix for high energies. *arXiv:1110.3710*, 2011.
[8] V. Guillemin. Sojourn times and asymptotic properties of the scattering matrix. In *Proceedings of the Oji Seminar on Algebraic Analysis and the RIMS Symposium on Algebraic Analysis (Kyoto Univ., Kyoto, 1976)*, volume 12, pages 69–88, 1976/77 supplement.
[9] A. Hassell and J. Wunsch. The semiclassical resolvent and the propagator for non-trapping scattering metrics. *Adv. Math.*, 217(2):586–682, 2008.
[10] L. Hörmander. Fourier integral operators. I. *Acta Math.*, 127(1-2):79–183, 1971.
[11] L. Hörmander. *The analysis of linear partial differential operators. I.* Classics in Mathematics. Springer-Verlag, Berlin, 1983.
[12] L. Hörmander. *The analysis of linear partial differential operators. III.* Classics in Mathematics. Springer-Verlag, Berlin, 2007. Pseudo-differential Operators, Reprint of the 1994 edition.
[13] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences.* Wiley-Interscience [John Wiley & Sons], New York, 1974. Pure and Applied Mathematics.
[14] A. Majda. High frequency asymptotics for the scattering matrix and the inverse problem of acoustical scattering. *Comm. Pure Appl. Math.*, 29(3):261–291, 1976.
[15] R. Melrose. *Geometric Scattering Theory.* Cambridge University Press, Cambridge, 1995.
[16] R. Melrose and M. Zworski. Scattering metrics and geodesic flow at infinity. *Invent. Math.*, 124(1-3):389–436, 1996.
[17] M. Reed and B. Simon. *Methods of modern mathematical physics. III.* Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1979.
[18] D. Yafaev. *Mathematical Scattering Theory: analytic theory.* American Mathematical Society, Providence, RI, 2010.
[19] D. R. Yafaev. On the asymptotics of scattering phases for the schrdinger equation. *Ann. Inst. H. Poincaré Phys. Théor.*, 53(3):283–299, 1990.
[20] S. Zelditch. Index and dynamics of quantized contact transformations. *Ann. Inst. Fourier (Grenoble)*, 47(1):305–363, 1997.
[21] S. Zelditch and M. Zworski. Spacing between phase shifts in a simple scattering problem. *Comm. Math. Phys.*, 204(3):709–729, 1999.

Department of Mathematics, Massachusettes Institute of Technology
E-mail address: `datchev@math.mit.edu`

Mathematical Sciences Institute, Australian National University and Department of Mathematics, University of Toronto
E-mail address: `jgell@math.toronto.edu`

Mathematical Sciences Institute, Australian National University
E-mail address: `Andrew.Hassell@anu.edu.au`

Mathematical Sciences Institute, Australian National University and Department of Mathematics, Princeton University
E-mail address: `peterch@math.princeton.edu`